

Nasal Air Flow During Normal Speech Production

AMY E. THOMPSON, M.S.
THOMAS J. HIXON, Ph.D.

Tucson, Arizona 85721

Nasal air flow was measured during the speech of 112 normal subjects (59 females and 53 males) ranging in age from three years to 37 years, six months. Flow was zero during nearly all oral consonant and vowel utterances, suggesting that velopharyngeal closure was air-tight. Flow occurred during all nasal consonants and during vowels adjacent to nasal consonants. Both age and sex effects were demonstrated for flow on vowels preceding nasal consonants. These effects were interpreted as showing that progressively older subjects and female subjects demonstrate earlier anticipatory coarticulation in preparation for forthcoming nasal consonants.

Introduction

The velopharyngeal mechanism is a mechanical valve situated between the oral and nasal portions of the upper airway. Its control of oral-nasal coupling accomplishes two important things in normal speech production: one, it enables the development of oral air pressure and air flow sufficient to produce many of the consonants of speech, and two, it enables the generation of voice perceived to be free of hypernasal quality. Despite the recognized importance of normal velopharyngeal function, many of the details of such function remain to be specified. The present study is one of a series of studies being conducted in the Department of Speech and Hearing Sciences at the University of Arizona to provide needed information about velopharyngeal function in normal speech production.

Three questions are of central importance in this part of this series of studies. The first concerns the extent to which air-tight velopharyngeal closure is characteristic of so-called phonetically-oralized speech elements. The impetus for this question is the belief that is commonly held by many individuals that complete closure of the velopharyngeal port

is not to be expected in normal speakers (Nusbaum, Foley, and Wells, 1935; Moll, 1962; Van den Berg, 1962).

The second question inquires into the possibility of a developmental schedule for gaining control over the velopharyngeal-closure mechanism. The impetus for this question is the lack of information pertinent to this topic and the fact that several motor-speech behaviors continue to be refined toward adultlike standards well after normal speech is acquired and sometimes as late as 11 to 12 years of age (Kent, 1976).

The third question of importance to the present investigation is an inquiry as to the possible existence of a sex difference with respect to control of the velopharyngeal-closure mechanism. A paucity of information exists in this regard, although that which is available (McKerns and Bzoch, 1970) would lead one to believe that there could be differences in function between the sexes based on differences in velopharyngeal structure.

The general design of the present investigation involved an attempt to address these three questions through the measurement of nasal air flow during the speech of a large group of males and females whose ages were distributed over a wide range. More specifically, subjects were selected to encompass a range extended downward to the youngest age at which standard tasks could easily be performed and extended upward into middle-aged adulthood. Flow was selected as the method for studying velopharyngeal function

The authors are affiliated with the University of Arizona, Tucson, Arizona. Dr. Hixon is a Professor in the Department of Speech and Hearing Sciences.

This work was supported by Research Grant NS-09656 from the National Institute of Neurological and Communicative Disorders and Stroke.

because of the simplicity of its measurement and because, for the purposes of answering the questions mentioned above, it constituted a sufficient tool for the study of normal speakers (Warren, 1967).

Method

SUBJECTS. One hundred twelve individuals served as subjects. Ninety-two of these subjects, 49 females and 43 males, ranged in age from three to 18 years. Ages were distributed relatively evenly so that at least one subject, and usually more than one, fell within each four-month interval within this age range. Typically, subjects of both sexes were included within each interval. The remaining persons studied were 20 adults, ten of each sex, ranging in age from 18 years, two months, to 37 years, six months. Most of these subjects were less than 25 years of age. All 112 subjects were selected to meet the following criteria: normal speech and voice commensurate with age, hearing reported to be within normal limits, no known structural or neurological disorders, and no known respiratory infections or allergies at the time of testing. All subjects were monolingual English speakers living in Tucson, Arizona.

EQUIPMENT. Two aspects of speech production were recorded: nasal air flow (hereinafter referred to as flow) and the acoustic pressure wave (hereinafter referred to as the speech signal). Flow was channeled through an individually-custom-formed mask placed over the nose. Masks were made of Dux-seal, a relatively compliant, clay-like material. Flow from the mask was sensed by a double-coned, two-square inch, Silverman-type, pneumotachometer coupled to a differential air-pressure transducer. The output analog voltage of the latter was amplified, and the resulting signal was low-pass filtered through a system with a sharp roll-off that markedly attenuated the signal above 20 Hz. The conditioned-flow analog was displayed on one channel of a two-channel storage oscilloscope. The flow analog was calibrated through use of a continuously-variable flow pump whose output was directed through a rotameter and the pneumotachometer in series. The speech signal was sensed by a condenser microphone positioned approximately six inches below and to the front of the subject's lips. The output from the microphone was amplified

and the resulting analog of the speech signal was displayed on the second channel of the storage oscilloscope. When it was deemed necessary to record data permanently, displays on the screen of the oscilloscope were photographed with a Polaroid camera. For certain activities, an electronic metronome was used to aid subjects in the pacing of their speech.

SPEECH SAMPLE. Utterance tasks consisted of three groups of activities: sustained productions, syllable repetitions, and nonsense productions embedded in a carrier phrase. Sustained productions included individual utterances of /i/, /s/, /z/, and /n/ at normal conversational pitch, loudness, and quality. Each utterance was made following an inspiration of about twice normal depth and lasted for about five seconds. Syllable repetitions included productions of /ti/, /di/, /si/, /zi/ and /ni/, each in trains of seven utterances, made at normal conversational pitch, loudness, and quality, with equal stress on each syllable, and at an utterance rate of three per second. Nonsense productions included the vowel-consonant-vowel (VCV) combinations /iti/, /idi/, /isi/, /izi/, and /ini/ in the carrier phrase "Say _____ again," produced at conversational pitch, loudness, and quality, with normal prosody, and with stress placed on the second vowel in the VCV. Each of the sustained productions and syllable repetition trains were performed three times in succession and each of the VCV productions was performed in a separate carrier phrase three times in succession. Utterances included in the study were chosen to encompass a variety of phonetic contexts that could be executed easily by the younger subjects. Design of the sample was also guided by a desire to include a variety of manners of consonant production (plosive, fricative, and nasal), and to include voicing contrasts for cognate pairs. A single place of consonant and vowel production was chosen, namely, lingua-alveolar. This place had the advantage of avoiding lip gestures that would interfere with the seal formed by the nasal mask. The high vowel /i/ was selected for study within the front vowel series because of its presumed stringent velopharyngeal-closure requirements (Moll, 1962), a factor we considered important for insuring the strength of a target behavior were a developmental schedule to be manifested. The utterance rate of three per second was found in a

pilot study to be the maximum rate at which very young subjects could perform reliably.

PROCEDURE. Each subject was seated upright and positioned so that the screen of the oscilloscope was out of the visual field. A mask was molded for the nose and was positioned on the subject. Each mask formed a small tube, the proximal end of which fitted airtight against the upper lip and mid-facial region surrounding the nose. Care was taken to insure that the walls of the mask did not block the external nares or contact the nasal alae. Next, the microphone was positioned, and the subject was instructed to repeat each utterance produced by an investigator. The model utterances for the subjects met the criteria discussed in the previous section. In the case of syllable repetitions, the investigator modelled utterance rate to the flashing and clicking of the metronome. Subject utterances judged to be inappropriate copies of the model were not accepted, and additional speech samples were elicited until an appropriate copy was obtained. Those utterances produced appropriately were stored on the screen of the oscilloscope for analysis. An investigator recorded the flow magnitudes of interest directly from the stored display. In cases where the flow analog could not be segmented instantly with regard to phonetic events, the oscilloscopic display was photographed for later perusal.

Results

We choose to divide the description of the results in accordance with the three groups of activities performed by the subjects: sustained productions, syllable repetitions, and nonsense productions embedded in a carrier phrase.

SUSTAINED PRODUCTIONS. The phonemes /i/, /s/, /z/, and /n/ comprised the group of sustained productions investigated. No nasal flow occurred for any of the utterances of the first three of these phonemes. That is, velopharyngeal closure presumably was air-tight for the 999 samples (3 phonemes $\times$ 3 repetitions $\times$ 111 subjects¹) involving sustained oral consonant or vowel elements. By contrast,

all 333 sustained productions of /n/ involved nasal flow. Criterion scores were computed for each of the subjects as the mean peak flow on the three /n/-productions. The resulting scores for the subject group yielded a mean value of 98.9 cc/sec, with a range of 23.3 cc/sec to 246.7 cc/sec, and a standard deviation of 38.1 cc/sec. Perusal of the criterion score data for /n/ revealed no indication that nasal flow was related either to the age or to the sex of the subjects studied.

SYLLABLE REPETITIONS. The syllables /ti/, /di/, /si/, /zi/, and /ni/, each performed in trains of seven, comprised the group of syllable repetitions. The middle three repetitions in each train were selected for analysis. No nasal flow occurred for any of the utterances of the syllables /ti/, /di/, /si/, and /zi/. That is, no flow occurred for 2664 sound elements (4 different syllables $\times$ 2 phonemes each $\times$ 3 repetitions $\times$ 111 subjects) involving oral consonant and vowel elements.² By contrast, of the 666 speech sounds - - 333 /n/'s and 333 /i/'s - - involved in the /ni/ syllable repetitions, all of the nasal consonants and all but nine of the vowels were accompanied by nasal flow. The nine speech sound utterances not accompanied by flow were performed by three subjects who uttered /i/ (three times each) without nasal flow. Criterion scores were computed for both /n/ and /i/ for each subject on the three productions of each phoneme. These computations were derived from

111 participants. To prevent this subject's data from unduly distorting the descriptive statistics for the overall group, we deemed it reasonable to isolate these data for separate discussion. They are considered in a subsequent footnote at the end of the RESULTS section.

² The statement "no nasal flow" should be qualified in the context of these first four syllable repetitions and in the context of the first four nonsense syllables studied to mean that no air passed from the mouth to the nose by way of the velopharynx. Very small magnitudes of nasal flow were observed for some subjects' productions of the utterances mentioned. These were typically less than ± 5 cc/sec, involved very gradual changes in flow, and often involved swings in flow from positive to negative and vice versa. Such observations are consistent with those of Lubker and Moll (1965), who demonstrated that their occurrence is related to minor displacements of the velum which alter the volume of the nasal cavity, change its internal pressure slightly, and cause small flows in and out of the nose in the presence of air-tight velopharyngeal closure. From the perspective of the measurements of importance to this investigation, these flows were viewed as artifacts and were specified as zero.

¹ Data for our subject group proved to be homogeneous. However, one of the 112 subjects presented data that usually were distinctly different from those of the other

measurements of flow made at the midpoint of each phoneme as determined from the accompanying speech signal. Flows measured at these points in the utterances proved to correspond closely to the maximum flow during the /n/ and the minimum flow during the /i/. Resulting scores were characterized by a subject group mean nasal flow for /n/ of 88.8 cc/sec, with a range of 40.0 cc/sec to 273.0 cc/sec, and a standard deviation of 34.1 cc/sec. For /i/ the subject group mean was 47.5 cc/sec, with a range of 0.0 cc/sec to 167.0 cc/sec, and a standard deviation of 25.3 cc/sec. Perusal of the data for /n/ and /i/ for the /ni/ utterances revealed no age or sex effects.

NONSENSE PRODUCTIONS EMBEDDED IN A CARRIER PHRASE. The syllables /iti/, /idi/, /isi/, /izi/, and /ini/, embedded in a carrier phrase, comprised the group of nonsense productions investigated. Nasal flow was zero for all of the 3996 individual speech sounds produced for the first four of these VCV syllables (4 syllables $\times$ 3 phonemes each $\times$ 3 repetitions $\times$ 111 subjects). The fifth of the nonsense syllables studied, /ini/, invariably was accompanied by nasal flow during the nasal consonant segment /n/. At the midpoint of /n/, flow for the subject group (for 333 utterances) was of a mean magnitude of 94.2 cc/sec, with a range of 30.0 cc/sec to 260.0 cc/sec, and a standard deviation of 38.9 cc/sec. Criterion score data for /n/ showed no indication of age or sex effects.

Flows measured at the midpoint of the initial vowel in the /ini/ nonsense productions resulted in a subject group mean value of 12.2 cc/sec, with a range of 00.0 cc/sec to 80.0 cc/sec, and a standard deviation of 15.9 cc/sec. Flows were zero at the midpoint of the initial vowel in /ini/ for 64 of the 111 subjects studied. Excluding these 64 subjects, recalculation of the mean, range, and standard deviation for the subject group resulted in values of 28.7 cc/sec, 3.3 cc/sec to 80.0 cc/sec, and 17.4 cc/sec, respectively. Criterion scores on the initial /i/ for our 111 subjects revealed two clear performance effects, one related to subject age and the other related to subject sex. With regard to age, there existed a moderate trend for an increasing proportion of the subjects to demonstrate nasal flow with increasing age. Considering the contiguous ages

from three to 18 years, and subdividing this range into three-year intervals, the data showed the following proportions of subjects to demonstrate nasal flow at the midpoint of the vowel in each interval, respectively: 13.6% at three to six years; 20.0% at six to nine; 42.9% at 9 to 12; 78.6% at 12 to 15; and 71.4% at 15 to 18 years. For the adult group, which covered a wide range of unevenly distributed ages, the proportion was 50%.

With regard to sex, 32 of the 47 subjects demonstrating nasal flow at the midpoint of the initial /i/ were female. This proportion (i.e., 68.1%) was observed not only overall but also in approximately the same magnitude across the five contiguous age subdivisions just mentioned and in the adult subject group.

For the final vowel in /ini/, flows measured at the midpoint yielded a subject group mean value of 43.3 cc/sec, with a range of 8.0 cc/sec to 120.0 cc/sec, and a standard deviation of 23.1 cc/sec. Examination of the criterion score data failed to reveal nasal flow effects for either age or sex.³

Discussion

The discussion of the results of this investigation will be organized under three subheadings: fully oralized utterances, fully nasalized utterances and simultaneously oralized and nasalized utterances, and clinical implications and research needs.

FULLY ORALIZED UTTERANCES. The fully oralized utterances of this investigation turned out to be the isolated utterances of /i/, /s/, and /z/, the syllable repetitions of /ti/, /di/, /si/, and /zi/, and the carrier phrase embedded syllables /iti/, /idi/, /isi/, and /izi/. Nasal flow was zero and the velopharynx presumably was closed air-tight for

³ As mentioned in Footnote 1, one of the subjects presented data that were single in kind for the majority of utterances sampled. This subject, a female, 13 years, seven months of age, performed in a manner typical of the overall subject group on sustained productions and the utterances of /ni/ and /ini/. However, she demonstrated flow on certain phonemes in non-nasal phonetic contexts for the syllable repetition and nonsense production activities. Nasal flow was found to be zero for all vowel utterances while a clear burst of flow occurred during the production of almost every /t/, /d/, /s/, and /z/. These bursts were of short duration and ranged in magnitude from 20 cc/sec to 80 cc/sec. Flow for the voiced element in each cognate pair was roughly twice that for the corresponding voiceless element.

all of these utterances for 111 of the subjects. The phonemes included in the fully oralized category are routinely categorized by phoneticians as oral consonants and vowels. The present findings indicate that these speech sounds are fully oralized by both sexes as early as age three and that the use of air-tight velopharyngeal closure for their production is maintained into middle-aged adulthood (37 years, 6 months, was the upper limit studied here).

These findings on fully oralized utterances are in agreement with those of other researchers who have used pneumotachometers to study nasal flow during speech samples similar to those used in the present investigation (Warren and DuBois, 1964; Lubker and Moll, 1965; Machida, 1967; Lubker, Schweiger, and Morris, 1970; Lubker, 1973). That is, oral consonants and vowels have been found by these other authors to be produced with no nasal flow or with very minimal flow. In those instances where minimal flow occurred, it appeared not to be the result of a velopharyngeal leak but of velar elevation in the presence of an air-tight velopharyngeal seal (Lubker and Moll, 1965; Lubker, 1973). Recall from footnote 2 that, in the present study, we treated such occurrences of flow as artifacts and specified them as zero.

Researchers who have used warm-wire flowmeters to study nasal flow during speech samples similar to ours, have provided data that are both supporting and conflicting with regard to our findings. For example, Quigley et al. (1963) and Quigley (1967) found essentially no nasal flow on utterances that are routinely classified as oral consonants and vowels. By contrast, both Van Hattum and Worth (1967) and Emanuel and Counihan (1970) report nasal flows for certain oral consonant and vowel elements. Both pairs of the latter authors acknowledge the fact that some of these flows are probably attributable to the occurrence of velar elevations in the presence of an air-tight velopharyngeal seal. In other cases, where the flows are seemingly of too large a magnitude to be accounted for solely by velar displacements, both pairs of authors express concern that these may reflect leaks between the oral and nasal sections of their flow-measuring apparatus. It is difficult to view either of those reports with much con-

viction because of the measurement problems they acknowledge. Hardy (1967) has severely criticized the instrumentation system used in the study by Van Hattum and Worth (1967) on a number of bases. Some of the same criticisms apply equally to the instrumentation of Emanuel and Counihan (1970).

Also relevant to the present discussion are available cinefluorographic data on velopharyngeal function during speech conditions of the nature studied in the present investigation. Moll (1962) presents data for velopharyngeal closure on vowels that reveal that, although the velopharynx usually is closed for vowel productions, it is open for a small percentage of utterances, usually in the neighborhood of 15% for the /i/-vowel contexts he studied. Bzoch (1968), by contrast, found no velopharyngeal opening in a cinefluorographic study of upper airway function. Aside from his formal investigation, Bzoch also inspected some 200 cinefluorographic films of normal speakers and found only occasional velopharyngeal opening on oral speech elements. He attributed these occurrences of velopharyngeal opening to slow utterance rates. We are inclined to accept Bzoch's conclusion based on our own further observations of several adult subjects who performed the study protocol at slower utterance rates than were studied in the 112 subjects. Two of six such subjects, all of whom demonstrated air-tight velopharyngeal closure for syllable repetitions at the usual rate of three per second, occasionally showed small nasal flows when utterance rate was dropped below 1.5 per second. Bzoch would appear to have made an important observation when he stated that "a slow rate of utterance of syllables is a factor related to the occasional occurrence of velopharyngeal opening on non-nasal speech elements." Clearly, the utterance rate factor needs to be specified or controlled in subsequent investigations.

Recall that one of the 112 subjects in this study failed to demonstrate air-tight velopharyngeal closure during some of the consonant elements in her productions of the syllable utterances of concern here. Her unique nasal flow patterns suggest that she briefly and slightly opened the velopharynx during the production of consonant elements but that she maintained air-tight velopharyngeal clo-

sure for adjacent vowel elements. Despite this idiosyncratic adjustment pattern, this subject's speech was normal in all regards. Although air-tight velopharyngeal closure is clearly typical for the oral consonants and vowels mentioned here, the data on this single subject with a velopharyngeal leak and normal speech raise the question about the necessity for air-tight closure. Other authors, most notably Warren (1967), have documented relatively small nasal flows in subjects judged to have adequate speech. For example, taking a very conservative velopharyngeal orifice size of 10 mm^2 as representing an "adequate" degree of velopharyngeal closure, one might expect to find nasal flows during voiceless stop-plosive productions that would reach magnitudes in the neighborhood of 100 cc per second. It is important to keep in mind that the amount of velopharyngeal opening that a listener will tolerate before labeling speech as deviant may depend upon whether the deviation pertains to hypernasal voice quality or to misarticulations - - the first predominantly an acoustics problem and the second predominantly an aeromechanics problem.

FULLY NASALIZED UTTERANCES AND SIMULTANEOUSLY ORALIZED AND NASALIZED UTTERANCES. We did not measure oral air flow in this investigation. Presumably, it was zero during the sustained productions of /n/, during the major part of the production time of all of the /n/'s in the syllable repetitions of /ni/, and in the syllable utterances of /ini/ embedded in the carrier phrase (Lubker and Moll, 1965; Subtelny, Worth, and Sakuda, 1966; Gilbert, 1968). Here, all /n/ productions are categorized as fully nasalized. The flow data for such nasalized utterances proved to be comparable for the three different phonetic contexts in which /n/ was studied. Furthermore, they revealed no apparent age or sex effects for this subject group. It is tempting to generalize these findings as indicative of fully matured velopharyngeal behavior that is acquired by at least age three and is maintained through middle-aged adulthood. It must be remembered, however, that several factors influence the flow of air from the nose during speech, including nasal pathway resistance, velopharyngeal airway resistance, and oral air pressure (Warren, 1967). The last of these alone, that is, oral pressure, is reported

to be influenced by factors such as phonetic context and age and sex of the subject, although the nature of such influences are not necessarily congruent from one study to another (Arkebauer, Hixon, and Hardy, 1966; Subtelny et al., 1966; Diggs, 1972; Bernthal and Beukelman, 1978). Too little is currently known to make much of the fact that the present data did not show effects for context, age, or sex.

The findings on fully nasalized utterances are in agreement with those of some authors (Warren and DuBois, 1964; Subtelny et al., 1966) but are in disagreement with those of others (Lubker and Moll, 1965; Lubker, et al., 1970). Specifically, in the present investigation, peak nasal flows were observed at magnitudes approaching about 100 cc per second on the average. Warren and DuBois (1964) and Subtelny et al., (1966) show data of these approximate magnitudes for utterances of /m/, /n/, and /ng/. By contrast, the studies of Lubker and Moll (1965) and Lubker et al. (1970), done in the same laboratory, present data that are typically twice the magnitude of those presented here and also by other investigators. Lubker and Moll (1965), for example, show flows of 200 cc per second peak magnitude for /n/ in running speech phrases comparable to ours, while Lubker et al. (1970) show flows of over 210 cc per second on the average for the /m/ in an /imi/ utterance context. It is difficult to reconcile flow magnitudes twice the magnitude of ours without invoking an explanation that embodies major laryngeal adjustments that would result in substantially higher flows being delivered to the oral cavity. We note that not only are the nasal flows reported by Lubker and his associates higher for the context mentioned above, but that they are also approximately double what we observed under conditions where flows from the nose were believed to be the result of velar adjustments in the presence of air-tight velopharyngeal closure. Perhaps the nasal-flow-channel calibration for the studies of Lubker and Moll (1965) and Lubker et al. (1970) was off by a constant factor related to the misreading of the standard scale. We have no other explanation.

The simultaneously oralized and nasalized utterances of this investigation were the /i/ vowels paired with the nasal consonant /n/

(that is, /ni/ and /ini/). Oralization is assumed for these utterances on the basis of past studies of oral air flow (Lubker and Moll, 1965; Subtelny et al., 1966; Machida, 1967; Van Hattum and Worth, 1967; Emanuel and Counihan, 1970). Evidence for the nasalized component of these utterances is clear from our nasal flow data and from the data of others (Warren and DuBois, 1964; Lubker and Moll, 1965; Van Hattum and Worth, 1967).

Our findings for nasal flow in the three different contexts for the /i/ vowel revealed that the /i/'s following the nasal consonant /n/ (that is, in /ni/ and in the second vowel in /ini/) were of closely comparable magnitudes while those measured on the /i/ preceding the nasal consonant were only about one-fourth the magnitude of the other two. The occurrence of nasal flow on /i/, a phoneme classified as oralized in typical phonetic schemes, is clearly attributable to the coarticulation of the nasal consonant /n/ with the vowels immediately preceding and following. In the case of the two contexts in which vowels follow the nasal, the occurrence of nasal flow on the vowel appears to be related to carry-over coarticulation in which the velopharynx does not close air-tight for the vowel after being open for the nasal. This finding is not unexpected and is consistent with the observations of others who have studied velopharyngeal function in similar contexts (Warren and DuBois, 1964; Lubker and Moll, 1965; Moll and Daniloff, 1971). Such carryover coarticulation is basically a consequence of the biomechanical nature of the speech mechanism.

For the utterance contexts in which /i/ preceded the nasal consonant (that is, the first vowel in /ini/), the occurrence of nasal flow on the vowel is clearly a result of another form of coarticulation, namely anticipatory. That is, the speaker begins to open the velopharynx during the initial vowel in /ini/ in anticipation of the nasal consonant to follow. This form of nasalization of the vowel in a nasal context has been observed by researchers using different observational techniques (Warren and DuBois, 1964; Moll and Daniloff, 1971).

With regard to age, our nasal flow data on the initial /i/ in /ini/ demonstrated a mod-

erate trend for a greater proportion of the subjects to show nasal flow at the mid-point of the initial vowel with increasing age. We interpret these data as evidence that progressively older subjects were demonstrating earlier anticipatory coarticulation in preparation for the forthcoming nasal. This progressively earlier occurrence of flow for older subjects constitutes a sort of "spread of nasalization" across the /ini/ syllable that moves in the direction of the first vowel. Unfortunately, our measurements at the mid-point of the vowel do not precisely define the function of the age effect we have mentioned. Obtaining the necessary information in this regard would require specification of the precise moment of velopharyngeal opening within each vowel utterance. Because data were not recorded permanently, such determinations cannot be made post-hoc. We are inclined to interpret our observations in terms of motor skills acquisition associated with the increased age of the subjects. That is, with increasing age, up to at least age 12, when behavior becomes more like that of adults, it appears that subjects take more and more advantage of preparatory gestures for following sounds. This may reflect an increased economy of biomechanics with increasing experience in using the speech mechanism. Kent (1976), for example, has summarized studies that show that several aspects of motor control for speech do not become fully adult-like until approximately this age. We believe the potential for understanding the developmental schedule for motor control of speech may be enhanced by more detailed studies of the spread of the nasalization phenomenon we have mentioned.

Recall that a clear sex effect was apparent in our flow data on the initial vowel in /ini/. Specifically, approximately 70% of the subjects who demonstrated nasal flow at the mid-point of the vowel were females, both overall and in the different age sub-groups. We are uncertain as to the reasons for this sex effect. Possibilities include a potentially greater tolerance for the spread of nasalization in females without the perception of hypernasality and biomechanical differences between the sexes that might favor greater anticipatory coarticulation by females. We lean toward the latter as a possible explanation for

our findings. McKerns and Bzoch (1970) have shown that velopharyngeal function differs considerably between males and females on a number of dimensions. Differences demonstrated included the orientation of the velum to the pharynx during velopharyngeal closure, the relationship of the inferior point of contact of the velum to the palatal plane, the degree of elevation of the velum, the amount of contact between the velum and the posterior pharyngeal wall, the distance from the uvula to the posterior pharyngeal wall, and the length of the velum in function for speech. It seems reasonable to suppose that other dynamic velopharyngeal function differences might exist between the sexes and that these might be manifested in measures such as the nasal-flow recordings made in the present study. That is, there may be certain mechanical advantages that would permit females and not males to engage in an earlier anticipatory opening of the velopharyngeal mechanism in preparation for /n/. Far too little is known about velopharyngeal biomechanics and aeromechanics to speculate with any degree of certainty in this regard.

CLINICAL IMPLICATIONS AND RESEARCH NEEDS. The results of this investigation appear to have implications for both clinical and research endeavors. It is our experience that many clinicians are of the opinion that some degree of velopharyngeal leakage may routinely accompany the production of what are phonetically classified as oralized speech sounds. The results of this investigation of a large number of normal subjects do not support this notion. Rather, it is apparent that, for the types of utterances sampled, air-tight velopharyngeal closure is nearly universal. This finding would appear to have implications with regard to the evaluation and the physical and behavioral management of persons with velopharyngeal incompetence. That is, the target for "normal" velopharyngeal closure is an air-tight seal. It remains to be determined what precisely the tolerable limits are for other than air-tight velopharyngeal closure. The fact that one of our subjects demonstrated normal speech in the presence of a small amount of nasal flow suggests that air-tight closure may not be a necessary requisite for normal speech. More detailed studies of subjects of this type may provide infor-

mation about the closure demands for normal speech production. Thus, the anticipated clinical observation in normal velopharyngeal closure on oral sounds would be no flow. However, a small amount of flow should not be interpreted to mean velopharyngeal incompetence. Fortunately, for clinical concerns, our results provide the first extensive data to illustrate that, at least for the oralized speech samples studied here, the clinician need not be concerned about subject differences with respect to age (over a wide range), sex, or phonetic context.

Our data on flow for the nasal consonant /n/ constitute a substantial body of information relative to the flow associated with fully nasalized consonants. They may well represent a useful standard of comparison for those concerned with secondary surgical procedures and prosthetic management techniques that occasionally "over-correct" for velopharyngeal incompetence to the extent that nasal breathing is impaired and hyponasality occurs. The central tendency and dispersion measures we have provided for nasal flow should provide a useful starting point for those interested in determining "typical" values for normal subjects.

With respect to research, there are clear needs to be filled. Knowing that air-tight velopharyngeal closure must develop sometime between birth and age three, it is obviously important to proceed with studies to determine the age at which children gain adult-like control of the velopharynx for oralized speech sounds. It should prove profitable to study such children with regard to both age and sex. New technological advances in flow measurement instrumentation, such as miniature thermistors, now make it possible to pursue studies of this type without encumbrance to the speech mechanism. Studies using such procedures are now underway in the University of Arizona Laboratories.

We acknowledge that the speech utterances studied in this investigation were limited with regard to sampling the phonemes of American English. Additional informal study in our laboratory of a small group of adults suggests to us that our findings of air-tight velopharyngeal closure on vowel utterances should not be generalized to all vowels. For example, some subjects demonstrate air-tight velophary-

ryngeal closure on high vowels but show small velopharyngeal leaks on low vowels. This being the case, subsequent research should take into account vowels in addition to the /i/ we studied here.

Finally, a number of aspects of velopharyngeal function cannot be specified with nasal-flow measurements alone. It would seem useful to undertake a comprehensive study of velopharyngeal function using other forms of innocuous aeromechanical procedures for quantifying such factors as velopharyngeal orifice area and the relationship between oral pressure and nasal flow. The technique of Warren and DuBois (1964) seems well suited to such pursuits.

Reprints: Thomas J. Hixon, Ph.D.
Professor

Department of Speech and Hearing Sciences
University of Arizona
Tucson, Arizona 85721

References

- ARKEBAUER, H., HIXON, T., AND HARDY, J., Peak intraoral air pressures during speech, *J. Speech Hear. Res.*, 10, 196-208, 1967.
- BERNTHAL, J., AND BEUKELMAN, D., Intraoral air pressure during the production of /p/ and /b/ by children, youths, and adults, *J. Speech Hear. Res.*, 21, 361-371, 1978.
- BZOCH, K., Variations in velopharyngeal valving: The factor of vowel changes. *Cleft Palate J.*, 5, 211-218, 1968.
- DIGGS, C., Intraoral air pressure for selected English consonants: A normative study of children, Master of Science Thesis, Purdue University, 1972.
- EMANUEL, F., AND COUNIHAN, D., Some characteristics of oral and nasal air flow during plosive consonant production, *Cleft Palate J.*, 7, 249-260, 1970.
- GILBERT, H., Oral airflow during stop consonant production, Master of Science Thesis, University of Wisconsin, 1968.
- HARDY, J., Techniques of measuring intraoral air pressure and rate of airflow, *J. Speech Hear. Res.*, 10, 650-654, 1967.
- KENT, R., Anatomical and neuromuscular maturation of the speech mechanism: Evidence from acoustic studies, *J. Speech Hear. Res.*, 19, 421-447, 1976.
- LUBKER, J., Transglottal airflow during stop consonant production. *J. Acoust. Soc. of Amer.*, 53, 212-215, 1973.
- LUBKER, J., AND MOLL, K., Simultaneous oral-nasal air flow measurements and cinefluorographic observations during speech production, *Cleft Palate J.*, 2, 257-272, 1965.
- LUBKER, J., SCHWEIGER, J., AND MORRIS, H., Nasal airflow characteristics during speech in prosthetically managed cleft palate speakers, *J. Speech Hear. Res.*, 13, 326-338, 1970.
- MACHIDA, J., Air flow rate and articulatory movement during speech, *Cleft Palate J.*, 4, 240-248, 1967.
- McKERNES, D., AND BZOCH, K., Variation in velopharyngeal valving: The factor of sex, *Cleft Palate J.*, 7, 652-662, 1970.
- MOLL, K., Velopharyngeal closure on vowels, *J. Speech Hear. Res.*, 5, 30-37, 1962.
- MOLL, K., AND DANILOFF, R., Investigation of the timing of velar movements during speech, *J. Acoust. Soc. of Amer.*, 50, 678-684, 1971.
- NUSBAUM, E., FOLEY, L., AND WELLS, C., Experimental studies of the firmness of the velar-pharyngeal occlusion during the production of English vowels, *Speech Monographs*, 2, 71-80, 1935.
- QUIGLEY, L., A comparison of air-flow and cephalometric techniques for evaluation of normal and cleft palate patients, *Amer. J. Orthod.*, 53, 423-445, 1967.
- QUIGLEY, L., WEBSTER, R., COFFEY, R., KELLEHER, R., AND GRANT, H., Velocity and volume measurements of nasal and oral air-flow in normal and cleft-palate speech, utilizing a warm-wire flow-meter and two-channel recorder, *J. Dent. Res.*, 42, 1520-1527, 1963.
- SUBTELNY, J., WORTH, J., AND SAKUDA, M., Intraoral pressure and rate of flow during speech, *J. Speech Hear. Res.*, 9, 498-518, 1966.
- VAN DEN BERG, J., Modern research in experimental phonetics, *Folia Phoniatrica*, 14, 81-149, 1962.
- VAN HATTUM, R., AND WORTH, J., Air flow rates in normal speakers, *Cleft Palate J.*, 4, 137-147, 1967.
- WARREN, D., Nasal emission of air and velopharyngeal function, *Cleft Palate J.*, 4, 148-156, 1967.
- WARREN, D., AND DUBOIS, A., A pressure-flow technique for measuring velopharyngeal orifice area during continuous speech, *Cleft Palate J.*, 1, 52-71, 1964.